

# FIB-4 Outperforms Liver Biopsy in the Assessment of Prognosis in HIV/HCV Coinfection

---

Berenguer J<sup>1</sup>, Zamora FX<sup>2</sup>, Carrero A<sup>1</sup>, von Wichmann MA<sup>3</sup>, López-Aldeguer J<sup>4</sup>, Téllez MJ<sup>5</sup>, Sanz J<sup>6</sup>, Esteban H<sup>7</sup>, Bellón JM<sup>1</sup>, González-García J<sup>2</sup>, and the GESIDA 3603 Study Group.

*<sup>1</sup>Hospital General Universitario Gregorio Marañón/IiSGM, <sup>2</sup>Hospital La Paz/IdiPAZ, <sup>3</sup>Hospital de Donostia, <sup>4</sup>Hospital La Fe, <sup>5</sup>Hospital Clínico San Carlos, <sup>6</sup>Hospital Príncipe de Asturias, <sup>7</sup>Fundación SEIMC/GESIDA, <sup>11</sup>Hosp. Germans Trias i Pujol, Badalona, SPAIN.*

**CROI 2014**  
**March 3-6, Boston, MA**  
**Presentation #: P-640**

# Background & Aims

---

- Liver biopsy (LB) is widely considered the gold standard in the assessment of the severity of liver disease <sup>1</sup>.
- FIB-4 is a non-invasive test (based on platelet count, age, AST, and INR) to estimate fibrosis in HIV/HCV-coinfected patients <sup>2</sup>.
- We compared the prognostic abilities of LB and FIB-4 in HIV/HCV coinfecting patients from the GeSIDA 3603 Study Cohort <sup>3</sup>.

1) Bravo AA, et al. **N Engl J Med** **2001**, 344: 495.

2) Sterling RK, et al. **Hepatology** **2006**; 43: 1317-25

3) Berenguer J, et al. **Clin Infect Dis** **2012**; 55(5): 728

# Methods

---

<b>Patients</b>	<ul style="list-style-type: none"><li>• From the GESIDA 3603 Study Cohort we selected patients with a baseline assessment of fibrosis by both LB (METAVIR) and FIB-4.</li></ul>
<b>Definitions of fibrosis</b>	<ul style="list-style-type: none"><li>• Significant fibrosis: LB <math>F \geq 2</math> / FIB-4 <math>\geq 1</math></li><li>• Advanced fibrosis: LB <math>F \geq 3</math> / FIB-4 <math>\geq 3.25</math></li></ul>
<b>Outcomes</b>	<ul style="list-style-type: none"><li>• Overall death</li><li>• Liver-related events: Decompensation or hepatocellular carcinoma (HCC), whichever occurred first.</li></ul>
<b>Statistics</b>	<ul style="list-style-type: none"><li>• We used ROC curves to determine the ability of LB and FIB-4 to predict outcomes. Comparisons were done by the method of Hanley and McNeil.</li><li>• We used adjusted Cox models to test the association of LB and FIB-4 with time to overall death.</li><li>• We used competing risk regression models to test the association of LB and FIB-4 with time to liver-related events (the competing risk was death).</li></ul>

# Results

---

<b>Patients</b>	<ul style="list-style-type: none"><li>• 903 patients (see <b>table 1</b>)</li></ul>
<b>Δt between LB and FIB-4</b>	<ul style="list-style-type: none"><li>• The median time (IQR) between LB and FIB-4 was 4.5 (2.2 – 12.0) months</li></ul>
<b>Follow-up</b>	<ul style="list-style-type: none"><li>• The median FU time was 63 months</li></ul>
<b>Events</b>	<ul style="list-style-type: none"><li>• Frequency of events during FU are shown in <b>table 2</b></li><li>• Accuracy of LB and FIB4 for the prediction of events are shown in <b>figure 1</b></li><li>• LB vs. FIB-4 cut-off points to rule-in &amp; rule-out events are shown in <b>table 3</b></li><li>• Kaplan-Meier estimates of liver-related events or overall death (whichever occurred first) for LB and FIB-4 are shown in <b>figure 2</b></li><li>• Adjusted hazard ratios (95% CI) of events according to LB and FIB-4 are shown in <b>figure 3</b></li></ul>

# Table 1. Baseline characteristics I

Characteristic	No SVR (n=575)	SVR (n=328)	Total (n=903)
Male sex *	437 (76.3)	241 (73.7)	678 (75.3)
Age – yr #	40 (37 - 43)	40 (37 - 43)	40 (37 - 43)
Prior IDU *	483 (84.7)	283 (86.8)	766 (85.5)
CDC category C *	139 (24.4)	68 (21.1)	207 (23.2)
CD4 + cells/uL nadir #	209 (104 - 322)	231 (140 - 353)	217 (116 - 330)
CD4 + cells/uL baseline#	532 (373 - 728)	527 (402 - 727)	529 (387 - 727)
HIV-RNA < LOQ *	366 (65.7)	213 (66.6)	579 (66)
Ethanol > 50 g/d*	30 (5.8)	8 (2.6) †	38 (4.6)
Methadone use	81 (15.2)	27 (8.8) †	108 (12.9)

\*n (%); # median (IQR)

† p<.05 with the group No SVR.

Abbreviations: IDU, injection drug use; LOQ, lower limit of quantification; LB, liver biopsy; Rx, treatment with IFN-RBV

# Table 1. cont. Baseline characteristics II

Characteristic	No SVR (n=575)	SVR (n=328)	Total (n=903)
HCV genotype*			
1 or 4	442 (76.9)	137 (41.8) †	579 (64.1)
2 or 3	118 (20.5)	183 (55.8) †	301 (33.3)
Unknown	15 (2.6)	8 (2.4)	23 (2.5)
HCV-RNA $\geq$ 500K IU/mL*	354 (73.4)	179 (61.1) †	533 (68.8)
METAVIR fibrosis score*			
F0, No. (%)	41 (7.1)	30 (9.1)	71 (7.9)
F1, No. (%)	137 (23.8)	105 (32) †	242 (26.8)
F2, No. (%)	147 (25.6)	89 (27.1)	236 (26.1)
F3, No. (%)	161 (28)	75 (22.9)	236 (26.1)
F4, No. (%)	89 (15.5)	29 (8.8) †	118 (13.1)
HBsAg positive*	26 (4.6)	8 (2.5)	34 (3.8)

\*n (%); # median (IQR)



† p<.05 with the group No SVR.

## Table 2. Frequency of events during follow-up

---

	<b>No SVR (N=575)</b>	<b>SVR (N=328)</b>	<b>Total (N=903)</b>	<b>P</b>
Overall deaths – n (%)	43 (7.5)	3 (0.9)	46 (5.1)	<.001
Liver decompensation – n (%)	61 (10.6)	3 (0.9)	64 (7.1)	<.001
Hepatocarcinoma – n (%)	11 (1.9)	2 (0.6)	13 (1.4)	.114
Liver-related event – n <sup>o</sup> (%)	67 (11.7)	4 (1.2)	71 (7.9)	<.001
Overall death/Liver-related event	83 (14.5)	7 (2.1)	90 (10)	<.001

**Fig 1. Accuracy of LB and FIB4 for the prediction of events**  
 Plotted = AUROCs (95% CI)



**Table 3. LB vs. FIB-4 cut-off points for advanced fibrosis to rule-in & rule-out events**

Liver-related events			
Cut-off	Event	No event	Total
<b>LB<math>\geq</math>F3</b>	45	309	354
<b>LB&lt;F3</b>	26	523	549
<b>Total</b>	71	832	903
<b>FIB4<math>\geq</math>3.25</b>	34	124	158
<b>FIB4&lt;3.25</b>	37	708	745
<b>Total</b>	71	832	903


Overall deaths/Liver-related events			
Cut-off	Event	No event	Total
<b>LB<math>\geq</math>F3</b>	54	300	354
<b>LB&lt;F3</b>	36	513	9
<b>Total</b>	90	813	903
<b>FIB4<math>\geq</math>3.25</b>	40	118	158
<b>FIB4&lt;3.25</b>	50	695	745
<b>Total</b>	90	813	903

	LB	FIB-4
<b>PPV % (95% CI)</b>	12.7 (9.1;16.3)	21.5 (14.8;28.2)
<b>NPV % (95% CI)</b>	95.3 (93.4;97.1)	95.0 (93.4;96.7)
<b>LR+ (95% CI)</b>	1.71 (1.40;2.08)	3.21 (2.40;4.30)
<b>LR- (95% CI)</b>	0.58 (0.43;0.80)	0.61 (0.49;0.77)
<b>Well classified</b>	63%*	82%*
*P<.001		

	LB	FIB-4
<b>PPV % (95% CI)</b>	15.2 (11.4;19.1)	25.3 (18.2;32.4)
<b>NPV % (95% CI)</b>	93.4 (91.3;95.6)	93.3 (91.4;95.1)
<b>LR+ (95% CI)</b>	1.63 (1.34;1.97)	3.06 (2.30;4.07)
<b>LR- (95% CI)</b>	0.63 (0.49;0.82)	0.65 (0.54;0.78)
<b>Well classified</b>	63%*	81%*
*P<.001		


# Figure 2. Kapla Meier estimates of Overall deaths/Liver-related events

## Liver biopsy




531	514	478	427	358	271	184	92	14
337	322	293	251	209	154	100	44	7

## FIB-4



720	701	653	581	487	361	240	118	19
148	135	118	97	80	64	44	18	2

**Figure 3. Adjusted HR (95% CI) of overall death and liver-related events according to absence or presence of advanced fibrosis assessed by both LB and FIB-4**



Adjusted by Age, Sex, HIV transmission category, CDC clinical category C, CD4+ cell nadir, HCV genotype, HCV RNA

# Conclusions

---

- ① We found that FIB-4 outperformed LB as a predictor of both overall death and liver-related events
- ② It must be taken into consideration that liver biopsy makes it possible to evaluate liver structure but not function, whereas FIB-4 enables us to evaluate liver function.
- ③ Our results call into question the role of liver biopsy as a gold standard for assessing prognosis in HIV/HCV coinfection.

# The GeSIDA 3603 Team

---

## Principal Investigators

J Berenguer, J Gonzalez

### H. Gregorio Marañón, Madrid

T Aldámiz, JM Bellón, J Cosín, I Gutiérrez, JC López, P Miralles, B Padilla, A Carrero, F Tejerina, C Diez, M Ramírez, M Sánchez-Conde, J Berenguer,

### H. 12 de Octubre, Madrid

MA Hernando, F Pulido, V Rodríguez, R Rubio

### H. Clinic, Barcelona

P Callau, JM Gatel, J Mallolas, JM Miro

### H. Clínico Univ de Valencia, Valencia

A Ferrer, MJ Galindo

### H. Clínico San Carlos, Madrid

MJ Téllez, J Vergas

### H. Donostia, San Sebastián

J Arrizabalaga, JA Iribarren, MA Von Wichmann

### H. General de Valencia, Valencia

E Ortega, L Ortiz

## Study Coordinators

E Barquilla, H Esteban

### H. La Paz, Madrid

J Alvarez, JR Arribas, I Bernardino, M Mora, F Pascual, JM Peña, E Rodríguez, I Valero, F Zamora, J González,

### H. Germans Trias i Pujol Badalona

B Clotet, A Jou, C Tural

### H. Getafe, Madrid

G Gaspar, G Pérez

### H. Guadalajara, Guadalajara

M Rodríguez, ML Montes

### H. La Fe, Valencia

S Cuellar, J López-Aldeguer

### H. La Princesa, Madrid

I Santos, J Sanz

### H. Móstoles, Madrid

C Barros, E Condés

## Statistician

JM Bellón

### Fund SEIMC-GESIDA, Madrid

E Aznar, E Barquilla, H Esteban, B Moyano

### H. Príncipe Asturias, Madrid

A Arranz, J de Miguel, J Sanz

### H. Ramón y Cajal, Madrid

A Moreno, S Moreno, C Quereda, MA Sanfrutos

### H. Santa Creu i Sant Pau Barcelona

P Domingo, JM Guardiola

### H. Severo Ochoa, Madrid

M Cervero, JJ Jusdado, R Torre

### H. Vall d'Hebron, Barcelona

M Crespo, E Van den Eynde

# Funding

---

- **FIPSE** (Refs. 36443/03, 36702/07, 361020/10)
- **FIS** (Refs. EC07/90734, PI11/01556, EC11/241)
- **RIS** (Ref RD12/0017)
- **I3SNS** (J Berenguer) (Refs. INT10/009, INT12/154)